arXiv:1409.4058v3 [math-ph] 28 Jan 2015

Commuting differential operators of rank 2 with polynomial coefficients.

V.Oganesyan.

Introduction.

If two differential operators

$$L_n = \sum_{i=0}^{n} u_i(x)\partial_x^i, \quad L_m = \sum_{i=0}^{m} v_i(x)\partial_x^i$$

commute, then there is a nonzero polynomial $R(z,w)$ such that $R(L_n, L_m) = 0$ (see [1]). The curve Γ defined by $R(z,w) = 0$ is called the *spectral curve*. If

$$L_n\psi = z\psi, \quad L_m\psi = w\psi,$$

then $(z,w) \in \Gamma$. For almost all $(z,w) \in \Gamma$ the dimension of the space of common eigenfunctions ψ is the same. The dimension of common eigenfunctions of two commuting differential operators is called the *rank*. The rank is a common divisor of m and n.
If the rank equals 1, then there are explicit formulas for coefficients of commutative operators in terms of Riemann theta-functions (see [2]).
The case when rank is greater than one is much more difficult. The first examples of commuting ordinary scalar differential operators of the nontrivial ranks 2 and 3 and the nontrivial genus g=1 were constructed by Dixmier [8] for the nonsingular elliptic spectral curve $w^2 = z^3 - \alpha$, where α is arbitrary nonzero constant:

$$L = (\partial_x^2 + x^3 + \alpha)^2 + 2x,$$

$$M = (\partial_x^2 + x^3 + \alpha)^3 + 3x\partial_x^2 + 3\partial_x + 3x(x^2 + \alpha),$$

where L and M is the commuting pair of the Dixmier operators of rank 2, genus 1. There is an example

$$L = (\partial_x^3 + x^2 + \alpha)^2 + 2\partial_x,$$

$$M = (\partial_x^3 + x^2 + \alpha)^3 + 3\partial_x^4 + 3(x^2 + \alpha)\partial_x + 3x,$$

where L and M is the commuting pair of the Dixmier operators of rank 3, genus 1.

The general classification of commuting ordinary differential operators of rank greater than 1 was obtained by Krichever [3]. The general form of commuting operators of rank 2 for an arbitrary elliptic spectral curve was found by Krichever and Novikov [4]. The general form of operators of rank 3 for an arbitrary elliptic spectral curve (the general commuting operators of rank 3, genus 1 are parametrized by two arbitrary functions) was found by Mokhov [5], [6]. Moreover, examples of commuting ordinary differential operators of arbitrary genus and arbitrary rank with polynomial coefficients were constructed in [11]. However, even in those cases for which explicit formulas were obtained the task of allocating commuting operators with polynomial coefficients is non-trivial and not completely solved until now. The task of a complete description of commuting differential operators with polynomial coefficients was posed and discussed by Mokhov in [10].
Mironov in [7] constructed examples of operators

$$L = (\partial_x^2 + A_3x^3 + A_2x^2 + A_1x + A_0)^2 + g(g+1)A_3x,$$

$$M^2 = L^{2g+1} + a_{2g}L^{2g} + ... + a_1L + a_0,$$

where a_i are some constants and A_i, $A_3 \neq 0$, are arbitrary constants. Operators L and M are commuting operators of rank 2, genus g.
Furthermore, in [9] it was proved by Mironov that the operators L_1 and M_1,

$$L_1 = (\partial_x^2 + \alpha_1\mathcal{P}(x) + \alpha_0)^2 + \alpha_1 g_2 g(g+1)\mathcal{P}(x), \quad \alpha_1 \neq 0,$$

$$M_1^2 = L_1^{2g+1} + a_{2g}L_1^{2g} + ... + a_1L_1 + a_0,$$

where a_i are some constants, α_i are arbitrary constants and $\mathcal{P}$ satisfies the equation

$$(\mathcal{P}'(x))^2 = g_2\mathcal{P}^2(x) + g_1\mathcal{P}(x) + g_0, \quad g_2 \neq 0,$$

where g_1 and g_2 are arbitrary constants, are a commuting pair of rank 2, genus g.
In the same paper it is proved that the operators L_2 and M_2,

$$L_2 = (\partial_x^2 + \alpha_1\wp(x) + \alpha_0)^2 + s_1\wp(x) + s_2\wp^2(x),$$

$$M_2^2 = L_2^{2g+1} + b_{2g}L_2^{2g} + ... + b_1L_2 + b_0,$$

where b_i are some constants, $\wp(x)$ is the Weierstrass elliptic function,
$\alpha_1 = \frac{1}{4} - 2g^2 - 2g$, $s_1 = \frac{1}{4}g(g+1)(16\alpha_0 + 5g_2)$, $s_2 = -4g(g+2)(g^2-1)$,

are also a commuting pair of rank 2, genus g.
In this paper we find new examples of commuting operators of rank 2. The following theorems are proved

Theorem 1. *Operator* $L = (\partial_x^2 + A_6x^6 + A_2x^2)^2 + 16g(g+1)A_6x^4$, *where* $g \in \mathbb{N}$,$A_6 \neq 0$ *and* A_2 *is an arbitrary constant, commutes with a differential operator* M *of order* $4m+2$, *for all* $m \geqslant g$. *The spectral curve has the form* $w^2 = z^{2m+1} + a_{2m}z^{2m} + ... + a_1z + a_0$. *The operators* L *and* M *are operators of rank 2.*

Theorem 2. *Operator* $L = (\partial_x^2 + A_4x^4 + A_2x^2 + A_0)^2 + 4g(g+1)A_4x^2$, *where* $g \in \mathbb{N}$, $A_4 \neq 0$ *and* A_2, A_0 *are arbitrary constants, commutes with a differential operator* M *of order* $4m+2$, *for all* $m \geqslant g$. *The spectral curve has the form* $w^2 = z^{2m+1} + a_{2m}z^{2m} + ... + a_1z + a_0$. *The operators* L *and* M *are operators of rank 2.*

Theorem 3.
1) *If* $L = (\partial_x^2 + A_nx^n + A_{n-1}x^{n-1} + ... + A_0)^2 + B_kx^k + B_{k-1}x^{k-1} + ... + B_0$, *where* $n > 3, n \in \mathbb{N}$, $A_n \neq 0$, $B_k \neq 0$, *commutes with an operator* M *of order* $4g+2$ *and* M, L *are operators of rank 2, then* $k = n-2$ *and* $B = (n-2)^2m(m+1)A_n$ *for some* $m \in \mathbb{N}$.
2) *For* $n > 6$ *operator* $L = (\partial_x^2 + A_nx^n)^2 + B_{n-2}x^{n-2}$ *does not commute with any differential operator* M *of order* $4g+2$, *where* M, L *could be a pair of rank 2.*
3) *If* $n = 5$, *then* $L = (\partial_x^2 + Ax^5)^2 + 18Ax^3$, $A \neq 0$, *commutes with an operator* M *of order* $4g+2$ *for all* g *and* M,L *are operators of rank 2.*
4) *Operator* $(\partial_x^2 + Ax^5)^2 + 9m(m+1)Ax^3$, $A \neq 0$ *does not commute with any differential operator* M *of order* $4g+2$, *for* $m > 1$, *where* M *and* L *could be a pair of rank 2* .

Note that for genus less than 9, for $m = g$, the spectral curve of the operator from theorem 1 is non-singular for almost all A_2. Apparently, when $m = g$, the spectral curve is non-singular for any g, but it is not proved. For $m > g$, apparently, the spectral curve is always singular. Here are few examples. For $m = g = 1$ the spectral curve of operator from theorem 1 has the form

$$w^2 = (z + 16A_2)(z^2 + 16A_2z + 192A_6).$$

For $g=1, m=3$, we have

$$w^2=(z+16A_2)(z^2+16A_2z+192A_6)(256A_2^2+C_2-16A_2C_1-zC_1(16A_2-1)+z^2)^2,$$

where C_1, C_2, C_3 are arbitrary constants. For $m=g=3$ and $A_2=0$

$$w^2=z(z^2+288000A_6)(273715200A_6^2+289152A_6z^2+z^4).$$

The operator from theorem 2 has singular spectral curve for $m=g=2,5$. Apparently the spectral curve is singular only for $m=g=3k-1$. Here are few examples. For $m=g=1$ the spectral curve has the form

$$w^2=z^3+8A_2z^2+16(A_2^2+A_0A_4)z+64A_0A_2A_4+16A_4^2.$$

For $g=1, m=3$ and $A_2=A_0=0$

$$w^2=(z^3+16A_4^2)(C_2+C_1z+z^2)^2,$$

where C_1, C_2 are arbitrary constants. If $g=m=3$ and $A_2=A_0=0$

$$w^2=z(3382560000A_4^4+117216A_4^2z^3+z^6).$$

Theorem 3 is not true for the general form of polynomials $Ax^n+A_{n-1}x^{n-1}+...+A_1x+A_0$ and $Bx^{n-2}+B_{n-1}x^{n-1}+...+B_1x+B_0$. For example, for any n there is a differential operator of order 6 that commutes with L. There is a reason to believe that L commutes with differential operator of order $4g+2$ for $g>1$ only for $n=3,4,5,6$, but it is not proved. Note that the spectral curve of operator $(\partial_x^2+Ax^5)^2+18Ax^3$ is always singular.

The author is grateful to O.I.Mokhov for valuable discussions.

Commuting operators of rank 2.

Consider the operator

$$(\partial_x^2+V(x))^2+W(x) \tag{1}$$

From article [7] we know that the operator commutes with an operator of order $4g+2$ with hyperlyptic spectral curve of genus g and hence is operator of rank 2, if and only if there is a polynomial

$$Q=z^g+a_1(x)z^{g-1}+a_2(x)z^{g-2}+...+a_{g-1}(x)z+a_g(x)$$

that the following relation is satisfied

$$Q^{(5)} + 4VQ''' + 6V'Q'' + 2Q'(2z - 2W + V'') - 2QW' \equiv 0, \qquad (2)$$

Q' means $\partial_x Q$. The spectral curve has the form

$$4w^2 = 4F(z) = 4(z-W)Q^2 - 4V(Q')^2 + (Q'')^2 - 2Q'Q''' + 2Q(2V'Q' + 4VQ'' + Q^{(4)}). \qquad (3)$$

Using (2), we get

$$4Q'z \equiv -Q^{(5)} - 4VQ''' - 6V'Q'' - 2Q'V'' + 2QW' + 4Q'W.$$

So we have the following system :

$$\begin{cases} a_1 = W/2 + C_1 \\ 4a_2' = -a_1^{(5)} - 4Va_1''' - 6V'a_1'' - 2a_1'V'' + 2a_1W' + 4a_1'W \\ \dots \\ 4a_{i+1}' = -a_i^{(5)} - 4Va_i''' - 6V'a_i'' - 2a_i'V'' + 2a_iW' + 4a_i'W \\ \dots \\ 4a_g' = -a_{g-1}^{(5)} - 4Va_{g-1}''' - 6V'a_{g-1}'' - 2a_{g-1}'V'' + 2a_{g-1}W' + 4a_{g-1}'W \\ 0 = -a_g^{(5)} - 4Va_g''' - 6V'a_g'' - 2a_g'V'' + 2a_gW' + 4a_g'W \end{cases}$$

It is clear from the system that result in [7] can be formulated in the following way. Let $a_1 = W/2 + C_1$, where C_1 is an arbitrary constant. Define a_i by recursion

$$a_{i+1} = \frac{1}{4}\int(-a_i^{(5)} - 4Va_i''' - 6V'a_i'' - 2a_i'V'' + 2a_iW' + 4a_i'W)dx \qquad (4)$$

We see that (1) commutes with an operator of order $4g+2$ and these operators are operators of rank 2 if and only if $a_{g+1} \equiv const$. For example,

$$a_2 = -\frac{1}{8}W^{(4)} - \frac{1}{2}VW'' - \frac{1}{4}V'W' + \frac{3}{8}W^2 + \frac{1}{2}WC_1 + 2C_2,$$

where C_2 is a constant, which appears after integration. In general, a_i contains i constants, i.e., $a_i(x) = a_i(x; C_1, ..., C_i)$.

Proof of Theorem 1.

Suppose that $V = A_6x^6 + A_2x^2$, $W = 16A_6g(g+1)x^4$, where $g \in \mathbb{N}$. The operator has the form

$$(\partial_x^2 + A_6x^6 + A_2x^2)^2 + 16A_6g(g+1)x^4. \tag{5}$$

Let us prove that we can choose constants $C_1, C_2, ..., C_g$ in such a way that $a_{g+1} = const$. Calculation shows that

$$a_1 = C_1 + 8A_6g(g+1)x^4.$$

It is clear from (4) that a_{i+1} is polynomial in x and linear in a_i. Let us apply (4) to x^{4k}. Assume that $a_i = x^{4k}$

$$a_{i+1} = C_{i+1} - k(4k-1)(4k-2)(4k-3)x^{4k-4} - 16A_2k^2x^{4k}$$
$$+\tfrac{8A_6}{k+1}(2k+1)(g-k)(g+k+1)x^{4k+4}.$$

Using this formula we can calculate the following

$$a_2 = \widetilde{C}_2 + 8A_6(\widetilde{C}_1 - 16A_2)g(g+1)x^4 + 96A_6^2g(g+1)(g-1)(g+2)x^8,$$

$$a_3 = \widetilde{C_3} + 8A_6g(g+1)(256A_2^2 - 16A_2\widetilde{C_1} + \widetilde{C_2} - 5040A_6(g-1)(g+2))x^4 -$$
$$-96A_6^2(80A_2 - \widetilde{C}_1)g(g+1)(g-1)(g+2)x^8 + 1280A_6^3g(g+1)(g-1)(g+2)(g-2)(g+3)x^{12},$$

where $\widetilde{C}_i$ are constants and depend on C_i. During the transition from a_3 to a_4 the term $\widetilde{C}_3$ will be multiplied by $8A_6g(g+1)x^4$. Also $16A_2 \cdot const \cdot x^4$ and the fourth derivation of $const \cdot x^8$ will be subtracted from $8\widetilde{C}_3A_6g(g+1)x^4$. Polynomial a_4 will have constant term $\widetilde{C}_4$.

Generally, if $a_i = \widetilde{C}_i + K_i^1x^4 + K_i^2x^8 + ... + K_i^ix^{4i}$, then

$$a_{i+1} = \widetilde{C}_{i+1} + (8A_6g(g+1)\widetilde{C}_i - 16A_2K_i^1 - 2 \cdot 7 \cdot 6 \cdot 5K_i^2)x^4 +$$

$$+(const \cdot K_i^1 - const \cdot K_i^2 - const \cdot K_i^3)x^8 + .. + \tfrac{8A_6(2i+1)}{i+1}(g-i)(g+i+1)K_i^ix^{4i+4}.$$

Also note that all coefficients, except the leading term, contain constants $C_1, C_2, ...$ And K_i^{i-1} contains only C_1, K_i^{i-2} contains only C_1 and C_2, K_i^{i-3} only C_1, C_2, C_3 etc.

We see that the term $const \cdot x^{4(g+1)}$ in a_{g+1} vanishes because it is multiplied by $const \cdot (g-g)(g+g+1) = 0$. Therefore, we must choose the constants $C_1, C_2, ...$ to vanish $K^m_{g+1} = 0$ for all m. This is always possible because the leading term in a_{g+1} equals $const \cdot x^{4g}$ and depends only on C_1, penultimate on C_1 and C_2 etc.
Let us suppose we didn't choose constants to vanish a'_{g+1} and continued applying transformation (4). Consider a_{g+1+m}, $m > 0$. Note that a_{g+1+m} has the same structure as a_{g+1} with the only difference that a_{g+1+m} contains constants $C_1, C_2, ..., C_{g+m}, C_{g+1+m}$. This means that we can choose constants $C_1, ..., C_{g+1+m}$ in such a way that $a'_{g+1+m} \equiv 0$. Hence there is a differential operator M of order $4(g+m)+2$ such that M commutes with L and they are operators of rank 2.

Proof of Theorem 2.

The operator has the form

$$(\partial_x^2 + A_4x^4 + A_2x^2 + A_0)^2 + 4g(g+1)A_4x^2. \tag{6}$$

As in the previous proof, we will prove that constants $C_1, C_2, ..., C_g$ can be chosen in such a way that $a_{g+1} = const$. Calculation shows that

$$a_1 = C_1 + 2A_4g(g+1)x^2.$$

We mentioned before that a_{i+1} is polynomial in x and linear in a_i. Let us apply (4) to x^{2k}. Assume that $a_i = x^{2k}$, then we have

$$a_{i+1} = C_{i+1} - k(2k-1)(k-1)(2k-3)x^{2k-4} - 2A_0k(2k-1)x^{2k-2} -$$

$$-4A_2k^2x^{2k} + \tfrac{2A_4(2k+1)(g-k)(g+k+1)x^{2k+2}}{k+1}.$$

Using this formula we can calculate the following

$$a_2 = \widetilde{C_2} + 2A_4(\widetilde{C_1} - 4A_2)g(g+1)x^2 + 6A_4^2g(g+1)(g-1)(g+2)x^4,$$

$$a_3 = \widetilde{C_3} + 2A_4g(g+1)(16A_2^2 - 4A_2\widetilde{C_1} + \widetilde{C_2} - 36A_0A_4(g-1)(g+2))x^2 -$$

$$-6A_4^2(20A_2 - \widetilde{C_1})g(g+1)(g-1)(g+2)x^4 + 20A_4^3g(g+1)(g-1)(g+2)(g-2)(g+3)x^6$$

where $\widetilde{C_i}$ are constants and depend on C_i.

Generally, let $a_i = \widetilde{C}_i + K_i^1 x^2 + K_i^2 x^4 + ... + K_i^i x^{2i}$, then

$a_{i+1} = \widetilde{C}_{i+1} + (2A_4 g(g+1)\widetilde{C}_i - 4A_2 K_i^1 - 12A_0 K_i^2 - 90K_i^3)x^2 +$

$+(const \cdot K_i^1 - const \cdot K_i^2 - const \cdot K_i^3 - const \cdot K_i^4)x^4 + ..+$

$+\frac{2A_4(2i+1)}{i+1}(g-i)(g+i+1)K_i^i x^{2i+2}.$

As in the proof of the previous theorem all terms except the leading term contain constants $C_1, C_2, ...$ And K_i^{i-1} contains only C_1, K_i^{i-2} contains only C_1, C_2 and K_i^{i-3} only C_1, C_2, C_3 etc.
The term $const \cdot x^{2(g+1)}$ in a_{g+1} vanishes because it is multiplied by $\frac{2A(2g+1)}{g+1}(g-g)(g+g+1) = 0$. So we must choose constants $C_1, C_2, ...$ in such a way that $A_{g+1}^i = 0$ for all i. This is always possible because the leading term in a_{g+1} depends only on C_1, penultimate on C_1 and C_2 etc.
Let us suppose we didn't choose constants to vanish a'_{g+1} and continued applying transformation (4). Consider a_{g+1+m}, $m > 0$. Note that a_{g+1+m} has the same structure as a_{g+1} with the only difference that a_{g+1+m} contains constants $C_1, C_2, ..., C_{g+m}, C_{g+1+m}$. This means that we can choose constants $C_1, ..., C_{g+1+m}$ in such a way that $a'_{g+1+m} \equiv 0$. Hence there is a differential operator M of order $4(g+m)+2$ such that L commutes with M.

Proof of Theorem 3.

If L commutes with operator of order $4g+2$, then $a_{g+1} \equiv 0$. We see that $a_i(x)$ is polynomial in x. Assume that degree of a_g equals l. Then degree of $-4Va_g''' - 6V'a_g'' - 2a_g'V''$ is equal to $n+l-3$. Sum $-4Va_i''' - 6V'a_i'' - 2a_i'V'' \neq 0$ because the leading coefficient is not equal to zero. Because of the same reason $2a_gW' + 4a_g'W \neq 0$. The degree of $2a_gW' + 4a_g'W$ equals $k+l-1$. So, if $a_{g+1} \equiv 0$, then $n+l-3 = k+l-1$, $\Rightarrow k = n-2$.
Let us prove that $B_{n-2} = (n-2)^2 m(m+1)A_n$ for some $m \in \mathbb{N}$. We will consider only leading coefficient of $a_i(x)$ and without loss of generality it can be assumed that the operator has the form

$$L = (\partial_x^2 + A_n x^n)^2 + B_{n-2}x^{n-2}, \tag{7}$$

We get

$$a_1 = C_1 + \frac{1}{2}B_{n-2}x^{n-2}.$$

We know that a_{i+1} is polynomial in x and linear in a_i. Let us apply (4) to x^k. Assuming that $a_i = x^k$

$$a_{i+1} = -\frac{1}{4}k(k-1)(k-2)(k-3)x^{k-4}+ \tag{8}$$

$$+\frac{(n+2k-2)}{2(n+k-2)}(B_{n-2} - Ak(n+k-2))x^{n+k-2} + C_{i+1}.$$

We see from this formula that the leading term of a_i is term of degree $(n-2)i$. If $a'_{g+1} \equiv 0$, then for some $m \leqslant g$ the leading term of a_{m+1} must vanish. Assuming in (4) that $a_i = x^{i(n-2)}$

$$a_{i+1} = -\frac{1}{4}i(n-2)(i(n-2)-1)(i(n-2)-2)(i(n-2)-3)x^{i(n-2)-4}+$$

$$+\frac{(2i+1)}{2(i+1)}(B_{n-2} - Ai(i+1)(n-2)^2)x^{i(n-2)+n-2} + C_{i+1}.$$

From this formula we have

$$B_{n-2} = (n-2)^2 m(m+1)A_n.$$

This means that we can rewrite (8) as

$$-\frac{1}{4}k(k-1)(k-2)(k-3)x^{k-4}+ \tag{9}$$

$$+\frac{A_n(n+2k-2)}{2(n+k-2)}(m(n-2)-k)((m+1)(n-2)+k)x^{n+k-2} + C_{i+1}.$$

Let us prove items 2–4. Assume $n > 6$.
Coefficient of the leading term in a_i is positive for $i < m$ because we see from (9) that during the transition from a_i to a_{i+1} the leading term will be multiplied by

$$\frac{A(2i+1)}{2(i+1)}(n-2)^2(m-i)(m+i+1)x^{n-2}.$$

By N_k denote $\frac{(n+2k-2)}{2(n+k-2)}(m(n-2)-k)((m+1)(n-2)+k)$.
As it was mentioned before $N_{i(n-2)} > 0$ for $i < m$. Easy to check that

$N_{i(n-2)-4} > 0$ for $i \leqslant m$, $i \geqslant 1$. Using (9), we get

$$a_2 = \widetilde{C}_2 - \frac{1}{8}Am(m+1)(n-5)(n-4)(n-3)(n-2)^3x^{n-6} + const \cdot x^{n-2} + \\ + \frac{3}{8}A^2(m-1)m(m+1)(m+2)(n-2)^4x^{2n-4}.$$

Note that the coefficient of the term x^{n-6} is negative. It is clear from (9) that during the transition to a_3 the term $A \cdot const \cdot x^{n-6}$ will be multiplied by $AN_{n-6}x^{n-2} = AN_{(n-2)-4}x^{n-2}$ and the fourth derivation of $A^2 \cdot const \cdot x^{2n-4}$ will be subtracted from $A \cdot const \cdot x^{2n-8}$. This means that the coefficient of x^{2n-8} in a_3 equals $A^2 \cdot const \cdot x^{2n-8}$, where constant is a negative number. So if $i \leqslant m$, then during the transition from a_i to a_{i+1} the coefficient of $x^{(i-1)(n-2)-4}$ will be multiplied by $AN_{(i-1)(n-2)-4}x^{n-2}$ and a positive number will be subtracted (the fourth derivation of $const \cdot x^{i(n-2)}$).
We see that if $i \leqslant m$, then the term $A^{i-1} \cdot const \cdot x^{(i-1)(n-2)-4}$ vanishes only if $A = 0$ because $const < 0$.
During the transition from a_m to a_{m+1} the leading term $const \cdot x^{m(n-2)}$ will be multiplied by 0 and will vanish.
If $i > m$, then during the transition from a_i to a_{i+1} nothing will be subtracted from the term $A^{i-1} \cdot const \cdot x^{(i-1)(n-2)-4}$. We see from (9) that during the transition to a_{i+1} the member $x^{(i-1)(n-2)-4}$ can vanish only because of factor $m(n-2) - (i-1)(n-2) + 4$. Suppose $i - m \geqslant 2$. Before we assumed that $n > 6$.

$$m(n-2) - (i-1)(n-2) + 4 = (n-2)(m-i+1) + 4 \leqslant -(n-2) + 4 < 0.$$

When $i - m = 1$

$$m(n-2) - (i-1)(n-2) + 4 = 4.$$

Finally, we obtain that if $n > 6$ and $A \neq 0$, then there is no such g that $a'_{g+1} \equiv 0$.

Assume $n = 5$. Consider operator

$$L = (\partial_x^2 + Ax^5)^2 + 18Ax^3. \tag{10}$$

Calculations show that

$$a_2 = \widetilde{C}_2 + 9A\widetilde{C}_1x^3.$$

So if $\widetilde{C}_1 = 0$, then $a_2' \equiv 0$. Hence, there is an operator M of order 6 such that M commutes with L.

$$a_3 = \widetilde{C}_3 + 9A\widetilde{C}_2 x^3$$

If $\widetilde{C}_2 = 0$, then there is an operator M of order 10 such that M commutes with L. In general

$$a_{g+1} = \widetilde{C}_{g+1} + 9A\widetilde{C}_g x^3.$$

If $\widetilde{C}_g = 0$, then there is an operator M of order $4g+2$ and M commutes with L.

Let $L = (\partial_x^2 + Ax^5)^2 + 9m(m+1)Ax^3$, where $m > 1$. Then

$$a_2 = \widetilde{C}_2 + 9\widetilde{C}_1 m(m+1)x^3 + \frac{243}{2}A^2 m(m+1)(m+2)(m-1)x^6$$

$$a_3 = \widetilde{C}_3 - \frac{10935}{4}A^2 m(m+1)(m+2)(m-1)x^2 + const\cdot x^3 + const\cdot x^6 + A^3\cdot const\cdot x^9$$

Arguing as before we will come to conclusion that the degree of x^2 will grow and never vanish.

References

[1] Burchnall J.-L., Chaundy T.W. Commutative ordinary differential operators. Proc. London Math. Soc. 21 (1923), 420-440; Proc. Royal Soc. London (A) 118 (1928), 557-583.

[2] I.M. Krichever, Integration of nonlinear equations by the methods of algebraic geometry, Func- tional Anal. Appl., 11: 1 (1977), 12–26.

[3] I.M. Krichever. Commutative rings of ordinary linear differential operators.Funktsional. Anal. i Prilozhen.,12:3 (1978), 20–31 (in Russian);English translation inFunctional Anal. Appl.,12:3 (1978), 175–185.

[4] I.M. Krichever, S.P. Novikov. Holomorphic bundles and nonlinear equations. Finite-zone solutions of rank 2.Dokl. Akad. Nauk SSSR,247:1 (1979), 33–36(in Russian);English translation in Soviet Math. Dokl..

[5] O.I. Mokhov. Commuting ordinary differential operators of rank 3 corresponding to an elliptic curve. Uspekhi Matem. Nauk,37:4 (1982), 169–170 (in Russian); English translation in Russian Math. Surveys,37:4 (1982), 129–130.

[6] O.I. Mokhov. Commuting differential operators of rank 3, and nonlinear differential equations.Izvestiya AN SSSR, ser. matem.,53:6 (1989), 1291–1315 (in Russian); English translation in Math. USSR, Izvestiya,35:3 (1990), 629–655.

[7] A.E. Mironov. Self-adjoint commuting differential operators and commutative subalgebras of the Weyl algebra. Invent. math. (2014) 197:417-431

[8] J. Dixmier,Bull. Soc. Math. France,96(1968), 209–24

[9] A.E.Mironov. Periodic and rapid decay rank two self-adjoint commuting differential operators arXiv:1302.5735

[10] O. I. Mokhov. On Commutative Subalgebras of the Weyl Algebra Related to Commuting Operators of Arbitrary Rank and Genus", Mat. Zametki, 94:2 (2013), 314–316

[11] O.I.Mokhov. Commuting ordinary differential operators of arbitrary genus and arbitrary rank with polynomial coefficients. American Mathematical Society Translations, Volume 234.

Department of Geometry and Topology, Faculty of Mechanics and Mathematics, Lomonosov Moscow State University, Moscow, 119991 Russia.

E-mail address: vardan.o@mail.ru